

This Page Is Inserted by IFW Operations
and is not a part of the Official Record.

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/33, 31/395, 31/40, 31/445, 31/495, 31/55, C07D 205/08, 207/12, 209/44, 211/40, 223/10, 225/02, 241/04, 295/08, 401/12, 403/12	A1	(11) International Publication Number: WO 98/51298 (43) International Publication Date: 19 November 1998 (19.11.98)
(21) International Application Number: PCT/US98/09023 (22) International Filing Date: 8 May 1998 (08.05.98) (30) Priority Data: 60/046,236 12 May 1997 (12.05.97) US (71) Applicant: ORTHO-McNEIL PHARMACEUTICAL INC. [US/US]; U.S. Route 202, Raritan, NJ 08869-0602 (US). (72) Inventors: JOLLIFFE, Linda; 16 Davenport Way, Belle Mead, NJ 08502 (US). MURRAY, William; 447 Township Line Road, Belle Mead, NJ 08502 (US). PULITO, Virginia; 37 Winding Way, Flemington, NJ 08822 (US). REITZ, Alan; 109 Greenbrian Road, Landsale, PA 19446 (US). LI, Xiaobing; 20 August Drive, Annadale, NJ 08801 (US). MULCAHY, Linda; 7 Jenny Jump Court, Flemington, NJ 08822 (US). MARYANOFF, Cynthia; 4029 Devonshire Drive, Forest Grove, PA 18922 (US). VILLANI, Frank; 2 Pinewood Lane, Perkasi, PA 18944 (US). (74) Agents: CIAMPORCERO, Audley, A., Jr. et al.; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08933 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: ARYLSUBSTITUTED PIPERAZINES USEFUL IN THE TREATMENT OF BENIGN PROSTATIC HYPERPLASIA		
(57) Abstract <p>This invention relates to a series of arylsubstituted piperazines, of Formula (I), pharmaceutical compositions containing them and intermediates used in their manufacture. The compounds of the invention selectively inhibit binding to the α-1a adrenergic receptor, a receptor which has been implicated in benign prostatic hyperplasia. As such the compounds are potentially useful in the treatment of this and other disease.</p> <div style="text-align: center;">  </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

ARYLSUBSTITUTED PIPERAZINES USEFUL IN THE TREATMENT OF BENIGN PROSTATIC HYPERPLASIA

This invention relates to a series of arylsubstituted piperazines,
5 pharmaceutical compositions containing them and intermediates used in their
manufacture. The compounds of the invention selectively inhibit binding to
the α -1_a adrenergic receptor, a receptor which has been implicated in benign
prostatic hyperplasia. In addition, compounds of the invention reduce
intraurethral pressure in an in vivo model. As such the compounds are useful
10 in the treatment of this disease.

BACKGROUND

Benign prostatic hyperplasia (BPH), a nonmalignant enlargement of
the prostate, is the most common benign tumor in men. Approximately 50%
15 of all men older than 65 years have some degree of BPH and a third of these
men have clinical symptoms consistent with bladder outlet obstruction (Hieble
and Caine, 1986). In the U.S., benign and malignant diseases of the prostate
are responsible for more surgery than diseases of any other organ in men
over the age of fifty.

20 There are two components of BPH, a static and a dynamic component.
The static component is due to enlargement of the prostate gland, which may
result in compression of the urethra and obstruction to the flow of urine from
the bladder. The dynamic component is due to increased smooth muscle
tone of the bladder neck and the prostate itself (which interferes with
25 emptying of the bladder) and is regulated by alpha 1 adrenergic receptors
(α 1-ARs). The medical treatments available for BPH address these
components to varying degrees, and the therapeutic choices are expanding.

Surgical treatment options address the static component of BPH and
include transurethral resection of the prostate (TURP), transurethral incision
30 of the prostate (TUIP), open prostatectomy, balloon dilatation, hyperthermia,
stents and laser ablation. TURP is the preferred treatment for patients with
BPH and approximately 320,000 TURPs were performed in the U.S. in 1990

at an estimated cost of \$2.2 billion (Weis et al., 1993). Although an effective treatment for most men with symptomatic BPH, approximately 20 - 25% of patients do not have a satisfactory long-term outcome (Lepor and Rigaud, 1990). Complications include retrograde ejaculation (70-75% of patients),
5 impotence (5-10%), postoperative urinary tract infection (5-10%), and some degree of urinary incontinence (2-4%) (Mebust et al., 1989). Furthermore, the rate of reoperation is approximately 15-20% in men evaluated for 10 years or longer (Wennberg et al., 1987).

Apart from surgical approaches, there are some drug therapies which
10 address the static component of this condition. Finasteride (Proscar[®], Merck), is one such therapy which is indicated for the treatment of symptomatic BPH. This drug is a competitive inhibitor of the enzyme 5 α -reductase which is responsible for the conversion of testosterone to dihydrotestosterone in the prostate gland (Gormley et al., 1992).
15 Dihydrotestosterone appears to be the major mitogen for prostate growth, and agents which inhibit 5 α -reductase reduce the size of the prostate and improve urine flow through the prostatic urethra. Although finasteride is a potent 5 α -reductase inhibitor and causes a marked decrease in serum and tissue concentrations of dihydrotestosterone, it is only moderately effective in
20 treating symptomatic BPH (Oesterling, 1995). The effects of finasteride take 6-12 months to become evident and for many men the clinical improvement is minimal (Barry, 1997).


The dynamic component of BPH has been addressed by the use of adrenergic receptor blocking agents (α 1-AR blockers) which act by
25 decreasing the smooth muscle tone within the prostate gland itself. A variety of α 1-AR blockers (terazosin, prazosin, and doxazosin) have been investigated for the treatment of symptomatic bladder outlet obstruction due to BPH, with terazosin (Hytrin[™], Abbott) being the most extensively studied. Although the α 1-AR blockers are well-tolerated, approximately 10-15% of
30 patients develop a clinically adverse event (Lepor, 1995). The undesirable effects of all members of this class are similar, with postural hypotension being the most commonly experienced side effect (Lepor et al., 1992). In

comparison to the 5 α -reductase inhibitors, the α 1-AR blocking agents have a more rapid onset of action (Steers, 1995). However, their therapeutic effect, as measured by improvement in the symptom score and the peak urinary flow rate, is moderate. (Oesterling, 1995)

- 5 The use of α 1-AR antagonists in the treatment of BPH is related to their ability to decrease the tone of prostatic smooth muscle, leading to relief of the obstructive symptoms. Adrenergic receptors are found throughout the body play a dominant role in the control of blood pressure, nasal congestion, prostrate function and other processes (Harrison et al., 1991). However,
- 10 there are a number of cloned α 1-AR receptor subtypes: α 1_a-AR, α 1_b-AR and α 1_d-AR (Bruno et al., 1991; Forray et al., 1994; Hirasawa et al., 1993; Ramarao et al., 1992; Schwinn et al., 1995; Weinberg et al., 1994). A number of labs have characterized the α 1-ARs in human prostate by functional, radioligand binding, and molecular biological techniques (Forray et al., 1994;
- 15 Hatano et al., 1994; Marshall et al., 1992; Marshall et al., 1995; Yamada et al., 1994). These studies provide evidence in support of the concept that the α 1_a-AR subtype comprises the majority of α 1-ARs in human prostatic smooth muscle and mediates contraction in this tissue. These findings suggest that the development of a subtype-selective α 1_a-AR antagonist might result in a
- 20 therapeutically effective agent with greater selectivity for the treatment of BPH.

SUMMARY OF THE INVENTION

The invention relates to compounds of Formula I



5

wherein:

A is $(CH_2)_n$ where n is 1-6;

R₁ is C₁₋₆alkyl, phenyl,
substituted phenyl

10

where the phenyl substituents are independently selected
from one or more of the group consisting of C₁₋₅alkyl,
C₁₋₅alkoxy and halogen, phenylC₁₋₅alkyl, or
substituted phenylC₁₋₅alkyl

15


where the phenyl substituents are independently selected
from one or more of the group consisting of C₁₋₅alkyl,
C₁₋₅alkoxy and halogen;

R₂ is hydrogen, C₁₋₆alkyl, C₁₋₅alkenyl, C₁₋₅alkynyl, phenylC₁₋₅alkyl,
or substituted phenylC₁₋₅alkyl

20

where the phenyl substituents are independently selected
from one or more of the group consisting of C₁₋₅alkyl,
C₁₋₅alkoxy and halogen;

E is



where:

m is 1-5;

5

R₃ is hydrogen, C₁₋₆alkyl or oxygen,

where if R₃ is oxygen, the hashed line represents a bond and if R₃ is C₁₋₆alkyl the hashed line is absent;

R₄ oxygen, hydrogen, C₁₋₅alkyl, formyl, carboxy, C₁₋₅alkylcarbonyl, C₁₋₅alkoxycarbonyl, phenylC₁₋₅alkoxy, substituted phenylC₁₋₅alkoxy

10

where the phenyl substituents are independently selected from one or more of the group consisting of C₁₋₅alkyl, C₁₋₅alkoxy and halogen, amido, and substituted amido

15

where the nitrogen substituents are independently selected from one or more of the group consisting of hydrogen, C₁₋₅alkyl, C₁₋₅alkoxy and hydroxy,

where if R₄ is oxygen, the hashed line represents a bond and if R₄ is any other substituent, the hashed line is absent;

20

R₅ is hydrogen, C₁₋₅alkyl or taken together with R₆ to form a cyclohexane, cyclopentane or cyclopropane ring;


R₆ is hydrogen, C₁₋₅alkyl or taken together with R₅ to form a cyclohexane, cyclopentane or cyclopropane ring;

and pharmaceutically acceptable salts thereof.

These compounds are useful as adrenergic receptor modulating agents. The compounds of this invention selectively bind to the α_{1a} -AR receptor, modulate the activity of said receptor and are selective for prostate tissue over aortic tissue. As such, they represent a viable treatment for α_{1a} -AR receptor modulated disorders which include but are not limited to BPH.

In addition this invention contemplates pharmaceutical compositions containing compounds of Formula I, and methods of treating disorders mediated by α_{1a} -AR receptor with compounds of Formula I.

Aside from compounds of Formula I, this invention contemplates intermediate compounds of Formula II. These intermediates are useful in the preparation of compounds of Formula I and are as follows:



wherein:

A is $(CH_2)_n$ where n is 1-6;

R_1 is C_{2-6} alkyl, phenyl, substituted phenyl


where the phenyl substituents are independently selected from one or more of the group consisting of C_{1-5} alkyl, C_{1-5} alkoxy and halogen, phenyl C_{1-5} alkyl, or substituted phenyl C_{1-5} alkyl

where the phenyl substituents are independently selected from one or more of the group consisting of C_{1-5} alkyl, C_{1-5} alkoxy and halogen;

R_7 is hydrogen, BOC or CBZ.

Still further this invention contemplates compounds of Formula III. These compounds are useful as intermediates in the preparation of compounds of Formula I and are as follows.

5



wherein:

m is 1-5.

10


DETAILED DESCRIPTION OF THE INVENTION

The terms used in describing the invention are commonly used and known to those skilled in the art. "HBSS" refers to Hank's Balanced Salt Solution. "Independently" means that when there are more than one
 15 substituent, the substituents may be different. The term "alkyl" refers to straight, cyclic and branched-chain alkyl groups and "alkoxy" refers to O-alkyl where alkyl is as known α 1-AR antagonists. defined supra. "LDA" refers to lithium diisopropylamide, and "BOP" refers to bezotriazole-1-yl-oxy-tris-(dimethylamino)-phosphoniumhexafluorophosphate. "BOC" refers to
 20 t-butoxycarbonyl, "PyBroP" refers to bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, "CBZ" refers to benzyloxycarbonyl, and "Ts" refers to toluenesulfonyl. "DCC" refers to 1,3-dicyclohexylcarbodiimide, "DMAP" refers to 4-N,N-dimethylaminopyridine, "EDCI" refers to 1-(3-dimethylamionpropyl)-3-ethylcarbodiimide hydrochloride, and "HOBT" refers
 25 to 1-hydroxybenzotriazole hydrate. The symbol "Ph" refers to phenyl and "PHT" refers to phthalimido. The term "effective dose" refers to an amount of a compound of Formula I which binds to an/or antagonizes the activity of the α -1_a adrenergic receptor. In addition, the term "effective dose" refers to an amount of a compound of Formula I which reduces the symptoms of diseases
 30 associated with the α -1_a adrenergic receptor.

The compounds of the invention may be prepared by the following schemes, where some schemes produce more than one embodiment of the invention. In those cases, the choice of scheme is a matter of discretion which is within the capabilities of those of synthetic chemists.

- 5 Compounds of Formula I where R_1 is phenyl, R_2 is hydrogen, R_3 is oxygen, A is $(CH_2)_3$ and m is 4, may be prepared as illustrated by Scheme 1. In this scheme, the molecules of Formula I are prepared in a convergent synthesis where two halves of the molecule are assembled and ultimately coupled together.
- 10 The starting material for one half is a mono N-substituted piperazine of type 1a. Treatment of 1a with a mild base such as K_2CO_3 and an alkylating agent, such as acrylonitrile in an inert solvent such as MeOH at room temperature for 2-24 hours gives the nitrile 1b. This intermediate may be hydrogenated using Rainy Nickel as a catalyst to give the propylamine
- 15 intermediate 1c. The other half of the molecule is assembled using compounds of type 1c as starting materials. ϵ -Caprolactam is treated with a strong base such as NaH, in an inert solvent at 0 °C for about 30 minutes. The formed anion is treated with an alkylating agent such as *t*-butylbromoacetate in an inert solvent such as acetonitrile at room
- 20 temperature for two hours to 2 days to give the ester 1e. Hydrolysis of 1e with an acid such as trifluoroacetic acid at room temperature over 2-16 hours gives the acid 1f. Treatment of the amine 1c and the acid 1f with a peptide coupling agent such as BOP and a suitable amine such as DMAP at room temperature over 1-16 hours gives a compound of Formula I, 1g. Other
- 25 coupling agents may be substituted for BOP. Such agents include but are not limited to, PyBroP, EDCI, HOBT and the like. Suitable DMAP replacements include but are not limited to N-methylmorpholine, imidazole, DABCO and the like.
- 30 Scheme 1 may be used to prepare compounds aside from 1g. To prepare compounds where m is 1-5, known lactams of the appropriate ring size such as 2-azacyclooctanone, replace ϵ -caprolactam in Scheme 1.

Scheme 1




To prepare compounds where R_1 is other than phenoxy, 1a is replaced
 5 with known phenyl piperazine derivatives such as N-(2-*t*-butoxyphenyl)-
 piperazine. To prepare compounds where A is $(CH_2)_n$ and n is 1-6,
 acrylonitrile may be replaced with alkylating agents such as
 4-chlorobutyronitrile. Alternatively, intermediate 1c may be prepared by
 another route as illustrated in Scheme 2.

10 Piperazine derivative 1a is treated with a mild base and known
 phthalimido derivatives such as N-(4-bromobutyl)phthalimide to give
 intermediate 2a. Treatment of 2a with a hydrazine such as N-methyl
 hydrazine in a suitable solvent such as MeOH or EtOH at reflux, gives
 intermediates of type 1c. Alternatively, piperazine 1a is treated with a mild


base and an N-BOC protected amine such as N-tert-butoxycarbonyl-4-bromobutylamine to give the corresponding BOC protected amine. This amine can be deprotected by treatment with an acid such as TFA to give intermediates of type 1c.

5

Scheme 2

To prepare compounds where R_2 is other than hydrogen, Scheme 3 may be used. Treatment of compounds of type 1g with a strong base such as NaH, followed by an alkylating agent such as benzyl bromide gives compounds of type 3a in 1-5 hours at temperatures from about 0-35 °C.

10

Scheme 3


Compounds where R_4 is oxygen, C_{1-5} alkyl, formyl, carboxy, C_{1-5} alkylcarbonyl, C_{1-5} alkoxycarbonyl, phenyl C_{1-5} alkoxy, substituted phenyl C_{1-5} alkoxy, amido and substituted amido may be synthesized by Scheme 4. For example to prepare compounds where R_4 is ethoxycarbonyl, m is 2, and R_3 is carbonyl, treat ethyl-2-pyrrolidone-5-carboxylate with a strong base such as NaH, in an inert solvent at 0 °C for about 30 minutes.

15


20

The formed anion is treated with an alkylating agent such as *t*-butylbromoacetate in an inert solvent such as acetonitrile at room temperature for two hours to 2 days to give the ester 4a. Acidic hydrolysis of 4a with trifluoroacetic acid at room temperature over 2-16 hours gives the acid 4b. Coupling of the acid 4b and the intermediate amine 1c with a peptide coupling agent such as PryBOP gives a compound of Formula I, 4c. The ethyl ester of compound 4c may be treated with a variety of agents to give derivatives such as amides, carboxylic acids, aldehydes etc, where the reagents and reaction conditions are within the knowledge of those skilled in the art.

Scheme 4




Although the claimed compounds are useful as modulators of $\alpha 1$ -AR, some compounds are either preferred or particularly preferred. The preferred
15 compounds of the invention include:



The particularly preferred "R₁"s are C₁₋₅alkyl.

The particularly preferred "R₂"s are hydrogen, C₁₋₅alkenyl and C₁₋₅alkynyl.

10 The particularly preferred "E"s are



The particularly preferred "m" is 3.

The particularly preferred R₃ is oxygen.

The particularly preferred R₄ is hydrogen.

15 The particularly preferred "A"s are (CH₂)_n where n is 2-4.

The particularly preferred compounds of Formula II include compounds where A is 2 or 3, R₁ is C₂₋₆alkyl, and R₇ is hydrogen.

The particularly preferred compounds of Formula III include compounds where m is 3.

5 The compounds of Formula I may be used in pharmaceutical compositions to treat patients with disorders related to modulating the activity of the α ₁ adrenergic receptor. The preferred route is oral administration, however compounds of the invention may be administered by other methods, including but not limited to intravenous infusion. Oral doses range from about
10 0.01-100 mg/kg daily. The preferred oral dosage range is from about 0.05-1.0 mg/kg daily. Infusion doses can range from about 0.001-100 mg/kg/min of inhibitor, with a pharmaceutical carrier over a period ranging from several minutes to several days.

 The pharmaceutical compositions can be prepared using conventional
15 pharmaceutical excipients and compounding techniques. Oral dosage forms may be elixirs, syrups, capsules tablets and the like. Where the typical solid carrier is an inert substance such as lactose, starch, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, mannitol and the like; and typical liquid oral excipients include ethanol, glycerol, water and the like.
20 All excipients may be mixed as needed with disintegrants, diluents, granulating agents, lubricants, binders and the like using conventional techniques known to those skilled in the art of preparing dosage forms. Parenteral dosage forms may be prepared using pharmaceutically acceptable carriers or diluents, including but not limited to water or another sterile
25 carrier.


 Typically the compounds of Formula I are isolated and used as free bases, however the compounds may be isolated and used as their pharmaceutically acceptable salts. Examples of such salts include but are not limited to hydrobromic, hydroiodic, hydrochloric, perchloric, sulfuric,
30 maleic, fumaric, malic, tartatic, citric, benzoic, mandelic, methanesulfonic, hydroethanesulfonic, benzenesulfonic, oxalic, pamoic, 2-naphthalenesulfonic, p-toluenesulfonic, cyclohexanesulfamic and saccharinic.

In order to illustrate the invention the following examples are included. These examples do not limit the invention. They are meant only to suggest a method of practicing the invention. Those skilled in the art may find other methods of practicing the invention, which will be readily apparent to them.

5 However those methods are deemed to be within the scope of this invention.

PREPARATIVE EXAMPLES


Example 1



10 1-(2-phthalimidoethyl)-4-(2-isopropoxyphenyl)piperazine
Cpd. 1

N-(2-Bromoethyl)phthalimide (7.6 g, 30 mmol) and K₂CO₃ (6.2 g, 45 mmol) were added to a solution of N-1-(2-isopropoxyphenyl)piperazine (6.6 g, 30 mmol) in acetonitrile (100 mL) and the resulting mixture was heated
15 at reflux for 2 days. The mixture was concentrated in vacuo and purified by column chromatography on silica gel using EtOAc/hexanes (30:70) as an eluent gave the title compound as a solid: MS m/z 394 (MH⁺).

Example 2




20 1-(2-Aminoethyl)-4-(2-isopropoxyphenyl)piperazine
Cpd. 2

A solution of compound 1 (7.5 g, 19 mmol) in EtOH (70 mL) was stirred for 10 minutes at room temperature. Methylhydrazine (20 mL) was added
25 and the mixture was heated at reflux for 2.5 hours. The mixture was cooled to room temperature and the resulting solid precipitate was removed by

filtration. The filtrate was concentrated in vacuo yielded the title compound as solid which was used without purification: MS m/z 264 (MH⁺).


Example 3

1-*t*-Butoxycarbonylmethyl-2-piperidone

Cpd. 3

95% Sodium Hydride (1.67 g, 66 mmol) was added to a stirred solution of δ -valerolactam (5.95 g, 60 mmol) in toluene (100 mL) at 0 °C and the resulting suspension was stirred for 1 hours. *t*-Butylbromoacetate (8.86 mL, 60 mmol) was added dropwise and the reaction mixture was warmed to room temperature and stirred for 10 hours. Saturated NH₄Cl aq was added and the resulting organic layer was washed with successive portions of brine and H₂O. The combined organic layers were dried (Na₂SO₄) and concentrated in vacuo to give compound 3 as an oil.

Example 4




1-Carboxymethyl-2-piperidone

Cpd 4

Trifluoroacetic acid (15 mL) was added to a stirred solution of compound 2 (12.93 g, 61 mmol) in methylene chloride (30 mL) under N₂. This mixture was stirred for 4 hours and concentrated in vacuo to give the title compound as a solid: MS m/z 158 (MH⁺).

Example 5



N-[Ethyl-2-(2-iso-propyloxyphenyl)piperazin-4-yl]-
[1'-(2-oxy-piperdiny)]acetamidemide


5

Cpd. 5

A solution of compound 2 (3.61 g, 23 mmol) in methylene chloride (10 mL) was added to a solution of compound 4 (5.0 g, 19 mmol) in methylene chloride (10 mL). PyBroP (10.72 g, 23 mmol), DMAP (3.85 g, 31 mmol) and N-methylmorpholine (2.53 mL, 23 mmol) were added and the mixture was stirred at room temperature under N₂ for 10 hours. The resulting mixture was washed with one portion of aqueous sodium bicarbonate, brine and H₂O. The combined organic layer was dried (Na₂SO₄) and concentrated in vacuo. The residue was purified by MPLC on silica gel using EtOAc/MeOH/triethylamine (90:5:5) as an eluent to give the title compound as an oil: MS m/z 403 (MH⁺). Treatment of the isolated compound with an equimolar portion of citric acid in ethyl acetate gave the citrate salt of the title compound as a solid.

15

Example 6



20

N-[Ethyl-2-(2-iso-propyloxyphenyl)piperazin-4-yl]-**N**-methyl-
[1'-(2-oxy-piperdiny)]acetamidemide

Cpd. 6

Sodium hydride (95% tech. 10.6 mg, 0.44 mmol) was added to a stirred solution of compound 5 (140.5 mg, 0.35 mmol) in THF (5 mL) at 0 °C and the

25

resulting suspension was stirred for 30 minutes. Methyl iodide (0.37 mmol) was added dropwise and the mixture was stirred overnight at room temperature. The residue was concentrated in vacuo, dissolved in ethyl acetate and washed with successive portions of aqueous, sat. ammonium chloride solution, brine and
5 water. The combined organic layer was dried (Na_2SO_4) and concentrated in vacuo to give the title compound as a solid: MS m/z 417 (MH^+).

BIOLOGICAL EXAMPLES

Biological activity and selectivity of compounds of the invention was
10 demonstrated by the following in vitro assays. The first assay tested the ability of compounds of Formula I to bind to membrane bound receptors α_{1a} -AR, α_{1b} -AR and α_{1d} -AR.

Example 14

The DNA sequences of the three cloned human α_1 -AR subtypes have
15 been published. Furthermore, the cloned cDNAs have been expressed both transiently in COS cells and stably in a variety of mammalian cell lines (HeLa, LM(tk-), CHO, rat -1 fibroblast) and have been shown to retain radioligand binding activity and the ability to couple to phosphoinositide hydrolysis. We used published DNA sequence information to design primers for use in RT-
20 PCR amplification of each subtype to obtain cloned cDNAs. Human poly A+ RNA was obtained from commercially available sources and included hippocampus and prostate samples, sources which have been cited in the literature. For the primary screen a radio ligand binding assay was used which employed membrane preparations from cells expressing the individual
25 cloned receptor cDNAs. Radiolabeled ligands with binding activity on all three subtypes (non-selective) are commercially available ($[^{125}\text{I}]$ -HEAT, $[^3\text{H}]$ -prazosin).

Each α_1 receptor subtype was cloned from poly A+ RNA by the standard method of reverse transcription-polymerase chain reaction (RT-
30 PCR). The following sources of polyA+ RNA were used for the cloning of the α_1 receptor subtypes: α_{1a} -AR, human hippocampus and prostate, α_{1b} -AR, human hippocampus, α_{1d} -AR, human hippocampus. The resulting cDNAs


were cloned into the pcDNA3 mammalian expression vector (Invitrogen Corp., San Diego CA). Each DNA was sequenced for verification and to detect any possible mutations introduced during the amplification process. Any deviation in sequence from the published consensus for each receptor subtype was corrected by site-directed mutagenesis.

The three $\alpha 1$ -AR subtypes (a, b, d) were transfected into COS cells using a standard DEAE-dextran procedure with a chloroquine shock. In this procedure, each tissue culture dish (100mm) was inoculated with 3.5×10^6 cells and transfected with 10 μ g of DNA. Approximately 72 hours post-transfection, the cells were harvested and COS membranes were prepared. Transfected COS cells from 25 plates (100mm) were scraped and suspended in 15mL of TE buffer (50mM Tris-HCl, 5mM EDTA, pH7.4). The suspension was disrupted with a homogenizer. It was then centrifuged at 1000xg for 10 minutes at 4 °C. The supernatant was centrifuged at 34,500xg for 20 minutes at 4 °C. The pellet was resuspended in 5mL TNE buffer (50mM Tris-HCl, 5mM EDTA, 150mM NaCl, pH7.4). The resulting membrane preparation was aliquoted and stored at -70°C. The protein concentration was determined following membrane solubilization with TritonX-100.

The ability of each compound to bind to each of the $\alpha 1$ -AR subtypes was assessed in a receptor binding assay. [125I]-HEAT, a non-selective $\alpha 1$ -AR ligand, was used as the radiolabeled ligand. Each well of a 96-well plate received: 140 μ L TNE, 25 μ L [125I]-HEAT diluted in TNE (50,000 cpm; final concentration 50 pM), 10 μ L test compound diluted in DMSO (final concentration 1 pM-10 μ M), 25 mL COS cell membrane preparation expressing one of the three $\alpha 1$ -AR subtypes (0.05-0.2 mg membrane protein). The plate was incubated for 1 hour at room temperature and the reaction mixtures were filtered through a Packard GF/C Unifilter filter plate. The filter plate was dried for 1 hour in a vacuum oven. Scintillation fluid (25 mL) was added to each well, and the filter plate was counted in a Packard Topcount scintillation counter. Data was analyzed using GraphPad Prism software.

Tables A & B list the IC₅₀ values expressed in nanomolar concentration for select compounds of the invention in all receptor subtypes.

Table A




5

Cpd.	R ₁	A	R ₂	R ₄	m	α1a-AR	α1b-AR	α1d-AR
5*	i-prop	CH ₂	H	H	3	8.7	46120	372
6	i-prop	CH ₂	CH ₃	H	3	53	763900	650
7	i-prop	CH ₂	CH ₂ CHCH ₂	H	3	103	172500	372
10 8	i-prop	CH ₂	CH ₂ (C)CH ₃	H	3	237	170200	358
9	i-prop	CH ₂	CH ₂ Ph	H	3	88	4226	348
10	i-prop	CH ₂	H	CO ₂ C ₂ H ₅	2	20	26310	260
11	i-prop	CH ₂	H	H	4	18	3536	505
12	CH ₃	(CH ₂) ₂	H	H	3	98	109000	30530

15 *** indicates a citrate salt

Table B



20

Cpd.	R ₁	A	R ₂	m	α1a-AR	α1b-AR	α1d-AR
13	i-prop	CH ₂	H	3	18	1932	189

Example 15

The antagonist activity of select compounds of Formula I was demonstrated by the following screen. Binding of an agonist to α1-ARs

causes the activation of PLC through G-protein coupled mechanisms (Minneman and Esbenshade, 1994). PLC catalyzes the hydrolysis of phosphatidyl inositol 4,5-bisphosphate (PIP₂) generating two second messenger molecules, inositol 1,4,5-triphosphate (IP₃) and diacylglycerol (DAG) and ultimately resulting in mobilization of intracellular calcium stores. Hits from the primary screening assay that showed selectivity in inhibiting radioligand binding to the α _{1a}-AR were evaluated for the ability to antagonize the mobilization of cytosolic calcium in cell lines stably expressing individual receptor subtypes.

Preparation of stable cell lines expressing each receptor subtype: The alpha 1 adrenergic receptor subtypes (a, b, d) in the pcDNA3 vector were transfected into HEK 293s (human embryonic kidney) cells to form stable receptor-expressing cell lines. Transfection was performed using DMRIE-C (GibcoBRL) cationic lipid reagent mixed with 2-3 ug of DNA and reduced serum OPTI-MEM 1 medium (GibcoBRL). Cells in 100mm tissue culture plates were overlaid with the lipid-DNA complex and incubated at 37°C, 5% CO₂ for 5-6 hours. Serum-containing growth medium was then added and cells were incubated for an additional 48 hours. Following this incubation, each plate was split 1:5 into selection medium containing 250, 300 or 350 ug/ml of G418 (geneticin) antibiotic. Plates were fed every 4 days with the appropriate selection medium. After approximately 3 weeks, colonies were picked for each subtype from the 300 ug/ml G418 selection plates. Colonies were expanded and frozen. Twelve cell lines of each subtype were screened for alpha 1 adrenergic receptor binding using a whole cell receptor binding assay. Positive cultures were subsequently analyzed by the calcium mobilization assay.

HEK 293s cells expressing human α _{1a}-AR were lifted with trypsin and washed once with HBSS. The cell pellet was resuspended in HBSS with 0.05% BSA to 1-5x10⁶ cells/ml. A 5 mM Fluo-3 solution (in 2/3 vol. DMSO and 1/3 vol Pluronic acid) was added to the cell suspension, giving a final Fluo-3 concentration of 5 μ M. Cells were then incubated with gentle rocking at room temperature for 1 hr in the dark. After incubation, the cells were

washed 3x with HBSS and resuspended in HBSS with 1.25 mM CaCl_2 to 0.7×10^6 cell/ml. Cell aliquots (100 μl) were pipetted into each well of a 96-well microplate. Calcium mobilization was induced with norepinephrine (10 μM) at room temperature. Two minutes prior to assay, antagonists were
5 added to cells utilizing a 96-well pipettor. Afterward, the agonist was added and fluorescent signal was monitored for 2-3 minutes using FLIPR (Molecular Devices, USA). Compound 5 inhibited the mobilization of cytosolic calcium at an IC_{50} of 99 μM .


Example 16

10 The antagonist activity and the selectivity of compounds of the invention for prostate tissues over aortic tissues as well as their antagonists was demonstrated as follows. The contractile responses of rat prostatic tissue and rat aorta tissues were examined in the presence and absence of antagonist compounds. As an indication of the selectivity of antagonism, test
15 compound effects on vascular smooth muscle contractility (α_{1b} -AR and α_{1d} -AR) were compared to the effects on prostatic smooth muscle (α_{1a} -AR). Strips of prostatic tissue and aortic rings were obtained from Long Evans derived male rats weighing 275 grams and sacrificed by cervical dislocation. The prostate tissue was placed under 1 gram tension in a 10 ml bath
20 containing phosphate buffered saline pH 7.4 at 32 ° C and isometric tension was measured with force transducers. The aortic tissue was placed under 2 grams tension in a 10 ml bath containing phosphate buffered saline pH 7.4 at 37 °C. The ability of test compound to reduce the norepinephrine-induced contractile response by 50 % (IC_{50}) was determined. Compound 5 inhibited
25 the contractile response in aortic tissue with an IC_{50} of 31.9 μM and in prostate tissue with an IC_{50} of 1.3 μM .

Example 17

Select compounds of the invention were tested for their ability to antagonize phenylephrine (PE) induced increases in intraurethral pressure in
30 dogs. The selectivity of these compounds was demonstrated by comparing their effect upon PE induced increases in mean arterial pressure (MAP) in the dog.


Male beagle dogs were anesthetized and catheterized to measure intraurethral pressure (IUP) in the prostatic urethra. Mean arterial pressure (MAP) was measured using a catheter placed in the femoral artery. Dogs were initially administered six i.v. bolus doses (1 to ≤ 32 mg/kg) of phenylephrine (PE) to establish a control agonist dose-response curve. IUP and MAP were recorded following each dose until the IUP returned to baseline. The dogs then were given an i.v. bolus dose of the antagonist compound, followed by i.v. PE challenges of ascending doses, as in the control agonist dose-response curve. IUP and MAP measurements following each PE challenge were recorded. The antagonist compound was tested over a dose range of 3 to 300 μ g/kg in half-log increments. The interval between antagonist doses was at least 45 minutes and three experiments were performed per dose level for each test compound. The graphs below illustrates the mean percentage reductions in IUP and MAP for compounds 5 and 12 respectively.



Effects of Compound 5 upon IUP and MAP at 10 μ g/kg PE in dogs

20

■ IUP
▲ MAP



Effects of Compound 12 upon IUP and MAP at 10 µg/kg PE in dogs

■ IUP
▲ MAP

5

REFERENCES

- Breslin D, Fields DW, Chou T-C, Marion DN, Kane M, Vaughan ED, and Felsen D (1993) Investigative urology: medical management of benign prostatic hyperplasia: a canine model comparing the in vivo efficacy of alpha-1 adrenergic antagonists in the prostate. J. Urol. 149: 395-399.
- Bruno JF, Whittaker J, Song J, and Berelowitz M. (1991) Molecular cloning and sequencing of a cDNA encoding a human α 1A adrenergic receptor. Biochem. Biophys. Res. Commun. 179: 1485-1490.
- Bylund DB, Eikenberg DC, Hieble JP, Langer SZ, Lefkowitz RJ, Minneman KP, Molinoff PB, Ruffolo RR, and Trendelenburg U (1994) IV. International Union of Pharmacology nomenclature of adrenoceptors. Pharmacol. Rev. 46: 121-136.
- Carruthers SG (1994) Adverse effects of α 1-adrenergic blocking drugs. Drug Safety 11: 12-20.

- Faure C, Gouhier C, Langer SZ, and Graham D (1995) Quantification of α 1-adrenoceptor subtypes in human tissues by competitive RT-PCR analysis. *Biochem. Biophys. Res. Commun.* 213: 935-943.
- 5 Flavahan NA and VanHoutte PM (1986) α -Adrenoceptor classification in vascular smooth muscle. *Trends Pharmacol. Sci.* 7: 347-349.
- Ford APDW, Arredondo NF, Blue DR, Bonhaus DW, Jasper J Dava MS, Lesnick J, Pfister JR, Shieh IA, Vimont RL, Williams RJ, McNeal JE, Stamey
10 TA, and Clarke DE (1996) RS-17053 (N-[2-(2-Cyclopropylmethoxyphenoxy)ethyl]-5-chloro-a, a-dimethyl-1H-indole-3-ethanamine hydrochloride), a selective α 1A-adrenoceptor antagonist, displays low affinity for functional α 1-adrenoceptors in human prostate: Implications for adrenoceptor classification. *Mol. Pharmacol.* 49: 209-215.
- 15 Forray C, Bard JA, Wegzel JM, Chiu G, Shapiro E, Tang R, Lepor H, Hartig PR, Weinshank RL, Branchek TA, and Gluchowski C (1994) The α 1-adrenergic receptor that mediates smooth muscle contraction in human prostate has the pharmacological properties of the cloned human α 1c
20 subtype. *Mol. Pharmacol.* 45: 703-708.
- Gormley G, Stoner E, Bruskewitz RC et al. (1992) The effects of finasteride in men with benign prostatic hyperplasia. *N. Engl. J. Med.* 327: 1185-1191.
- 25 Hatano A Takahashi H, Tamaki M, Komeyama T, Koizumi T, and Takeda M (1994) Pharmacological evidence of distinct α 1-adrenoceptor subtypes mediating the contraction of human prostatic urethra and peripheral artery. *Br. J. Pharmacol.* 113: 723-728.
- 30 Harrison JK, Pearson WR, and Lynch KR (1991) Molecular characterization of α 1- and α 1-adrenoceptors. *Trends Pharmacol. Sci.* 12: 62-67.

Hieble JP and Caine M (1986) Etiology of benign prostatic hyperplasia and approaches to pharmacological management. Fed. Proc. 45: 2601.

5 Hirasawa A, Horie K, Tanaka T, Takagaki K, Murai M, Yano J, and Tsujimoto G (1993) Cloning, functional expression and tissue distribution of human cDNA for the α 1c-adrenergic receptor. Biochem. Biophys. Res. Commun. 195: 902-909.

10 Holck CM, Jones CHM, and Haeusler G (1983) Differential interactions of clonidine and methoxamine with postsynaptic α -adrenoceptors of rabbit main pulmonary artery. Cardiovasc. Res. 5: 240-248.

15 Lepor H and Rigaud G (1990) The efficacy of transurethral resection of the prostate in men with moderate symptoms of prostatism. J. Urol. 143: 533-537.

20 Lepor H, Auerbach S, Puras-Baez A et al. (1992) A randomized, placebo-controlled multi-center study of the efficacy and safety of terazosin in the treatment of benign prostatic hyperplasia. J. Urol. 148: 1467-1474.

Lepor H (1995) α -Blockade for benign prostatic hyperplasia (BPH) J. Clin. Endocrinol. Metab. 80: 750-753.

25 Marshall I, Burt RP, Andersson PO, Chapple CR, Greengrass PM, Johnson GI, and Wyllie MG (1992) Human α 1c-adrenoceptor: functional characterisation in prostate. Br. J. Pharmacol. 112: 59P.

30 Marshall I, Burt RP, and Chapple CR (1995) Noradrenaline contraction of human prostate by α 1A- (α 1C-)adrenoceptor subtype. Br. J. Pharmacol. 115: 781-786.

Mebust WK, Holtgrewe HL, Cockett ATK, and Peters PC (1989) Transurethral prostatectomy: immediate and postoperative complication: a cooperative study of 13 participating institutions evaluating 3,885 patients
J. Urol. ,141: 243-247.

5

Minneman KP, Han C and Abel PW (1988) Comparison of α 1-adrenergic receptor subtypes distinguished by chloroethylclonidine and WB4101. Mol. Pharmacol. 33: 509-514.

- 10 Minneman KP and Esbenshade TA (1994) α 1-Adrenergic receptor subtypes. Annu. Rev. Pharmacol. Toxicol. 34: 117-133.

Morrow AL and Creese I (1986) Characterization of alpha 1-adrenergic receptor subtypes in rat brain: A reevaluation of [3H]WB4101 and

- 15 [3H]prazosin binding. Mol. Pharmacol. 29: 321-330.

Muramatsu I (1992) A pharmacological perspective of α 1-adrenoceptors: subclassification and functional aspects, in α -Adrenoceptors (Fujiwara M, Sugimoto T, and Kogure K, eds.). Excerpta Medica Ltd., Tokyo, 193-202.

20

Muramatsu I, Oshita M, Ohmura T, Kigoshi S, Akino H, Gobara M, and Okada K (1994) Pharmacological characterization of α 1-adrenoceptor subtypes in the human prostate: functional and binding studies. Br. J. Urol. 74: 572-577.


- 25 Oesterling JE (1995) Benign prostatic hyperplasia. Medical and minimally invasive treatment options. N. Engl. J. Med. 332: 99-109.

Price DT, Lefkowitz RJ, Caron MG, Berkowitz D, and Schwinn DA (1994) Localization of mRNA for three distinct α 1-adrenergic receptor subtypes in
30 human tissues: implications for human α -adrenergic physiology. Mol. Pharmacol. 45: 171-175.

- Ramarao CS, Kincade Denker JM, Perez DM, Gaivin RJ, Riek RP, and Graham RM (1992) Genomic organization and expression of the human α 1B-adrenergic receptor. *J. Biol. Chem.* 267: 21936-21945.
- 5 Schwinn DA, Johnston GI, Page SO, Mosley MJ, Wilson KH, Worman NP, Campbell S, Fidock MD, Furness LM, Parry-Smith DJ, Peter B, and Bailey DS (1995) Cloning and pharmacological characterization of human alpha-1 adrenergic receptors: sequence corrections and direct comparison with other species homologues. *JPET* 272: 134-142.
- 10 Weinberg DH, Trivedi P, Tan CP, Mitra S, Perkins-Barrow A, Borkowski D, Strader CD, and Bayne M (1994) Cloning, expression and characterization of human α adrenergic receptors α 1A, α 1B, and α 1C. *Biochem. Biophys. Res. Commun.* 201: 1296-1304.
- 15 Weis KA, Epstein RS, Huse DM, Deverka PA and Oster G (1993) The costs of prostatectomy for benign prostatic hyperplasia. *Prostate* 22: 325-334.
- Wennberg JE, Roos N, Sola L, Schori A, and Jaffe R (1987) Use of claims.
20 data systems to evaluate health care outcomes: mortality and reoperation following prostatectomy. *JAMA* 257: 933-936.
- Yamada S, Tanaka C, Kimura R, and Kawabe K (1994) Alpha 1-adrenoceptors in human prostate: characterization and binding
25 characteristics of alpha 1-antagonists. *Life Sci.* 54: 1845-1854.

What is claimed is:

1. A compound of Formula I



5

wherein:

A is $(CH_2)_n$ where n is 1-6;

R₁ is C₁₋₆alkyl, phenyl,
substituted phenyl

10

where the phenyl substituents are independently selected
from one or more of the group consisting of C₁₋₅alkyl,
C₁₋₅alkoxy and halogen, phenylC₁₋₅alkyl, or
substituted phenylC₁₋₅alkyl

15


where the phenyl substituents are independently selected
from one or more of the group consisting of C₁₋₅alkyl,
C₁₋₅alkoxy and halogen;

R₂ is hydrogen, C₁₋₆alkyl, C₁₋₅alkenyl, C₁₋₅alkynyl, phenylC₁₋₅alkyl,
or substituted phenylC₁₋₅alkyl

20

where the phenyl substituents are independently selected
from one or more of the group consisting of C₁₋₅alkyl,
C₁₋₅alkoxy and halogen;

E is



where:

m is 1-5;

R₃ is hydrogen, C₁₋₆alkyl or oxygen,

where if R₃ is oxygen, the hashed line represents a bond and if R₃ is C₁₋₆alkyl the hashed line is absent;

R₄ oxygen, hydrogen, C₁₋₅alkyl, formyl, carboxy, C₁₋₅alkylcarbonyl, C₁₋₅alkoxycarbonyl, phenylC₁₋₅alkoxy, substituted phenylC₁₋₅alkoxy

where the phenyl substituents are independently selected from one or more of the group consisting of C₁₋₅alkyl, C₁₋₅alkoxy and halogen, amido, and substituted amido

where the nitrogen substituents are independently selected from one or more of the group consisting of hydrogen, C₁₋₅alkyl, C₁₋₅alkoxy and hydroxy,

where if R₄ is oxygen, the hashed line represents a bond and if R₄ is any other substituent, the hashed line is absent;

R₅ is hydrogen, C₁₋₅alkyl or taken together with R₆ to form a cyclohexane, cyclopentane or cyclopropane ring;


R₆ is hydrogen, C₁₋₅alkyl or taken together with R₅ to form a cyclohexane, cyclopentane or cyclopropane ring;

and pharmaceutically acceptable salts thereof.

2. The compounds of claim 1 where R_1 is C_{1-6} alkyl, n is 2-4, and R_2 is hydrogen or C_{1-6} alkyl, C_{1-6} alkenyl.


5

3. The compounds of claim 2



4. The compounds of claim 2 where R_3 is oxygen and R_4 is hydrogen, C_{1-5} alkoxycarbonyl, or oxygen.

10

5. The compounds of claim 4 where R_4 is hydrogen and E is 

6. The compounds of claim 7 where m is 2 to 5.

15


7. A compound and pharmaceutically acceptable salts thereof selected from the group consisting of N-[ethyl-2-(2-iso-propyloxyphenyl)piperazin-4-yl)]-[1'-(2-oxy-piperdiny)]acetamidide, N-[ethyl-2-(2-iso-propyloxyphenyl)piperazin-4-yl)]-N-methyl-[1'-(2-oxy-piperdiny)]acetamidide, and N-[propyl-3-(2-iso-propyloxyphenyl)piperazin-4-yl)]-[1'-(2-oxy-piperdiny)]acetamidide.

20

8. A compound N-[ethyl-2-(2-iso-propyloxyphenyl)piperazin-4-yl)]-[1'-(2-oxy-piperdiny)]acetamidide and pharmaceutically acceptable salts thereof.

25

9. A pharmaceutical composition comprising a compound according to claim 1 and a pharmaceutically acceptable carrier or diluent.
10. A pharmaceutical composition comprising a compound according to claim 6 and a pharmaceutically acceptable carrier or diluent.
11. A pharmaceutical composition comprising a compound according to claim 8 and a pharmaceutically acceptable carrier or diluent.
12. A method of treating a disease mediated by the α -1_a adrenergic receptor comprising administering a compound of claim 1 to a patient at an effective dose.
13. A method of treating a disease mediated by the α -1_a adrenergic receptor comprising administering a composition of claim 6 to a patient at an effective dose.
14. A method of treating a disease mediated by the α -1_a adrenergic receptor comprising administering a composition of claim 8 to a patient at an effective dose.
15. The method of claim 18 where the compound is administered orally and an effective dose is 0.01-100 mg/kg daily.
16. The method of claim 15 where the dose is 0.05-1.0 mg/kg daily.
17. A method of treating benign prostatic hyperplasia comprising administering an effective dose of a compound of Formula I.
18. A compound of Formula II



II

wherein:

A is $(\text{CH}_2)_n$ where n is 1-6;

R₁ is C₂₋₆alkyl, phenyl,

5 substituted phenyl

where the phenyl substituents are independently selected

from one or more of the group consisting of C₁₋₅alkyl,

C₁₋₅alkoxy and halogen, phenylC₁₋₅alkyl, or

substituted phenylC₁₋₅alkyl

10 where the phenyl substituents are independently selected

from one or more of the group consisting of C₁₋₅alkyl,

C₁₋₅alkoxy and halogen;

R₇ is hydrogen, BOC or CBZ.

19. The compounds of claim 23 where n is 2-4 and R₁ is C₂₋₆alkyl.

15

20. The compounds of claim 25 where R₇ is hydrogen.


21. A compound selected from the group consisting of 1-(2-aminoethyl)-4-

(2-2-iso-propyloxyphenyl)piperazine, 1-(3-aminopropyl)-4-(2-2-iso-

20 propyloxyphenyl)piperazine, and 1-(4-aminobutyl)-4-(2-2-iso-
propyloxyphenyl)piperazine.

22. A compound 1-(2-aminoethyl)-4-(2-2-iso-propyloxyphenyl)piperazine.

25 23. A compound of Formula III



III

wherein:

m is 1-5.

30

24. The compounds of claim 28 where m is 2-5.
25. A compound, 1-t-butoxycarbonylmethyl-2-piperidone.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/09023

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/183, 210, 212, 255, 315, 424; 540/356, 362, 451, 463, 481, 524, 531, 598; 544/360, 362, 372, 373, 392; 546/243; 548/551

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4,841,051 A (SHIOZAWA et al.) 20 June 1989, column 4, lines 35-40.	23-25
X --- Y	WO 84/04302 A1 (BYK GULDEN LOMBERG CHEMISCHE FABRIK GESELLSCHAFT MIT BESCHRANKTER HAFTUNG) 08 November 1984, pages 2, 3, 14-16 and 18.	23-25 ----- 23-27
Y	US 4,069,336 A (LANGE et al.) 17 January 1978, column 2, line 11.	28-29



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents.	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 31 JULY 1998	Date of mailing of the international search report 24 AUG 1998
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <i>Brenda Coleman</i> BRENDA COLEMAN Telephone No. (703) 308-1235

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/09023

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 6, 10, 13, 15-21 and 25
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

Generic claim 6 is dependent on species claim 7, and claims 10 and 13 are dependent on claim 6. Claim 15 is dependent on non-existing claim 18 and claim 21 is dependent on claim 15. Claim 25 is dependent on itself. Claim 16-20 do not exist.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/09023

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 31/33, 31/395, 31/40, 31/445, 31/495, 31/55; C07D 205/08, 207/12, 209/44, 211/40, 223/10, 225/02, 241/04, 295/08, 401/12, 403/12

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

514/183, 210, 212, 255, 315, 424; 540/356, 362, 451, 463, 481, 524, 531, 598; 544/360, 362, 372, 373, 392; 546/243; 548/551